

Online Research @ Cardiff

This is an Open Access document downloaded from ORCA, Cardiff University's institutional repository: <https://orca.cardiff.ac.uk/id/eprint/125795/>

This is the author's version of a work that was submitted to / accepted for publication.

Citation for final published version:

Rosignon, Alexandre and Bonifazi, Davide ORCID: <https://orcid.org/0000-0001-5717-0121> 2019. O-Annulation leading to five-, six-, and seven-membered cyclic diaryl ethers involving C-H cleavage. *Synthesis* 51 (19) , pp. 3588-3599. 10.1055/s-0037-1611892 file

Publishers page: <http://dx.doi.org/10.1055/s-0037-1611892>
<<http://dx.doi.org/10.1055/s-0037-1611892>>

Please note:

Changes made as a result of publishing processes such as copy-editing, formatting and page numbers may not be reflected in this version. For the definitive version of this publication, please refer to the published source. You are advised to consult the publisher's version if you wish to cite this paper.

This version is being made available in accordance with publisher policies.

See

<http://orca.cf.ac.uk/policies.html> for usage policies. Copyright and moral rights for publications made available in ORCA are retained by the copyright holders.




O-Annulation Leading to Five-, Six-, and Seven-Membered Cyclic Diaryl Ethers Involving C–H Cleavage

Alexandre Rossignon^{a,b}

Davide Bonifazi^b

^aDepartment of Chemistry, University of Namur, Rue de Bruxelles 61, Namur 5000, Belgium

^bSchool of Chemistry, Cardiff University, Main Building, Park Place, Cardiff CF10 3AT, Wales, UK
bonifazi@cardiff.ac.uk



Abstract Cyclic diaryl ethers are present in multiple natural compounds, organic pollutants as well as in π -conjugated organic molecular materials. This short review aims at overviewing the main synthetic advances in the O-annulation methods for preparing five-, six-, and seven-membered rings through C–H cleavage.

- 1 Introduction
- 2 Five-Membered Rings: The Dibenzofuran (DBF) Motif
 - 2.1 Palladium-Catalysed C–H Activation
 - 2.2 Copper-Catalysed C–H Activation
 - 2.3 Non-CH Activation Oxidant-Mediated Cyclisation
 - 2.4 Light-Mediated Cyclisation
 - 2.5 Acid-catalysed C–O Cleavage/C–O Formation
- 3 Six-Membered Rings: DBX, PXX, Xanthone, and Their Derivatives
 - 3.1 Dibenzoxanthene (DBX)
 - 3.2 *Peri*-Xanthenoxanthene (PXX)
 - 3.3 Xanthenes
 - 3.4 Miscellaneous
- 4 Seven-Membered Rings: Cularine
- 5 Conclusion

Key words cyclic diaryl ether, oxidative cyclisation, cycloetherification, C–H activation, C–O bond formation, O-annulation, pyrans, furans, cularine

1 Introduction

Cyclic diaryl ether motifs are found in a large variety of organic molecules, spanning from natural products to polycyclic aromatic hydrocarbon (PAHs) structures (Figure 1). The antibiotic drug Vancomycin is probably the most famous structure.^{1a} Other examples include Vicanicin, a lichen secondary metabolite displaying antitumor activity,^{1b} and Asterelin A, an antifungal drug (Figure 1, top).^{1c}



Alexandre Rossignon (left) was born in Namur (Belgium) in 1988. He received a Master's degree in chemistry from the University of Namur (UNamur) in 2013. Then, he started his Ph.D. under the supervision of Prof. D. Bonifazi at the University of Namur. In 2015, he moved to Cardiff University as a visiting student in the School of Chemistry. His research interests include the synthesis of O-doped polycyclic aromatic hydrocarbons for applications in electrochromic devices.

Davide Bonifazi (right) was born in Guastalla (Italy) in 1975. After obtaining the 'Laurea' in 'Industrial Chemistry' from the University of Parma working with Prof. Enrico Dalcanale, he joined the group of Prof. François Diederich as a Ph.D. student at the ETH Zürich (2000–2004). He was awarded the Silver Medallion of the ETH for his doctoral dissertation (2005). After a one-year postdoctoral fellowship with Prof. Maurizio Prato at the University of Trieste, he joined the same University as a research associate first, and then as a part-time Professor (2012–2015). In 2006, he joined the University of Namur (BE) as Junior Professor (2006–2011) and as Professor of Organic Chemistry (2012–2015). Since 2016 he is Chair Professor of Organic Supramolecular Chemistry in the School of Chemistry at Cardiff University (UK). His activities are focused on the creation of functional organic architectures in interdisciplinary projects through targeted organic synthesis, self-assembly, and self-organisation of organic architectures in solution and on surfaces, physical-organic studies, material- and bio-based design.

Concerning artificial cyclic ethers, the polychlorinated dibenzo-*p*-dioxins and polychlorinated dibenzofurans are certainly the most infamous examples, being in the 'Dirty Dozen' list of Persistent Organic Pollutants (POPs) (Figure 1, centre).² Cyclic ethers also constitute the core of O-doped

polycyclic aromatic hydrocarbons (PAHs), a class of electron-rich aromatics used to engineer organic semiconductors (Figure 1, bottom). For instance, derivatives of Pummerer's *per*xanthenoxanthene (PXX) have been used as organic semiconductors for engineering the first Sony's 'Rollable OTFT-driven OLED that can wrap around a Pencil'.³ 2*H*-Pyran-

based dithienopyran (DTP)⁴ and dibenzopyran (DBP)⁵ motifs were introduced as the electron-donating moiety in donor-acceptor (D-A) type conjugated copolymers to engineer solar cells with high power conversion efficiencies (PCEs). Molecular units based on 9-[3-(dibenzo[*b,d*]furanyl)phenyl]-9*H*-carbazole (DFPCz) scaffold have been used as organic phosphors in PHOLEDs.⁶





Figure 1 Examples of natural and artificial products featuring cyclic ether structural motifs

Very interesting *p*-type organic semiconductor based on O-doped quinoidal pentacenes and nonacenes were recently prepared following a cross-condensation route.⁷ At the synthetic planning level, cyclic diaryl ethers are usually prepared through an intramolecular coupling reaction between a phenol moiety and a pre-functionalized aryl substrate. In particular, two routes have been mostly exploited:

(i) transition-metal-catalysed C–O cross-coupling reactions (i.e., Buchwald–Hartwig,⁸ Ullmann etherification,⁹ and Chan–Lam–Evans¹⁰ coupling), and (ii) nucleophilic aromatic substitution reactions.^{11,12} Both synthetic approaches rely on the use of electrophilic aromatic halides or strong acidic conditions to ensure the 'activation' of the aryl substrate and control the regioselectivity of the addition reaction (Scheme 1).



Scheme 1 General methodologies for preparing diaryl ethers


Also, the electrophilic partner needs to be pre-functionalised at a given position with electron-withdrawing groups (EWGs) and undergoes metal-arene formation,¹³ di-aryl iodonium salt,¹⁴ and in situ formation of benzyne species¹⁵ to favour the relevant addition reaction (Scheme 1). Cross-condensation reaction promoted by strong acids is also a valuable route when the substrate is not acid-sensitive.¹⁶ The interested reader can refer to recent reviews tackling these synthetic routes.¹⁷ A more attractive route would involve the formation of a cyclic diaryl ether through direct C–H functionalisation of an aryl substrate with a giv-

en phenolic moiety. It is with this aim that in this review paper we describe the recent synthetic developments tackling the formation of cyclic diaryl ethers using an intramolecular etherification route involving a C–H cleavage reaction. This review will provide an exhaustive picture of the synthetic plans involving the specific reactions and substrates to give five-, six-, and seven-membered rings.

2 Five-Membered Rings: The Dibenzofuran (DBF) Motif

2.1 Palladium-Catalysed C–H Activation


In 2011, Liu and co-workers¹⁸ described the first transition-metal-catalysed cyclisation reaction involving a C–H activation/C–O bond formation to obtain dibenzofuran derivatives by aerobic oxidation starting from 2-phenylphenol **1**. Their method involves the use of Pd(OAc)₂, 1,3-bis(2,6-diisopropylphenyl)imidazole-2-ylidene (IPr), mesitylene carboxylate (MesCOONa), 4,5-diazafluoren-9-one as ancillary ligands, and K₂CO₃ (Scheme 2). The use of the anionic ligand MesCOONa serves as a proton shuttle to promote C–H activation and 4,5-diazafluoren-9-one to favour the aerobic oxidation of the Pd(0)-based species to active Pd(II) complexes. Substrates bearing electron-donating groups (EDGs: amine, ether, ketal, and silyl) as well as electron-withdrawing groups (EWGs: cyano, ketone, nitro, amide, ester, sulfonamide, fluoride, chloride, and trifluoromethyl) on both the aryl and phenol moieties were used. Kinetic isotope effect (KIE) investigations suggested that the rate determining step is the C–O reductive elimination. Moreover, the reaction occurs quantitatively and regioselectivity at the least sterically hindered C–H positions.




Scheme 2 Palladium-catalysed DBF formation reported by Liu and co-workers¹⁸

In parallel, Wei and Yoshikai¹⁹ developed a base-free method using Pd(OAc)₂ and 3-nitropyridine as the ligand (Scheme 3). Under these conditions, aerobic oxidation is inefficient and BzOOtBu was used as an inexpensive oxidant

for the regeneration of the active palladium species. Mixture of aromatic and Lewis-basic solvents, such as C₆F₆ and DMI (1,3-dimethyl-2-imidazolidinone) (3:2 ratio), gave the highest yield. With respect to Liu's Pd-catalysed cycloetherification reaction, in this case, KIE investigations suggest that the rate determining step is the C–H bond cleavage ($k_H/k_D = 1.9$). Building on these kinetic studies, and considering that BzOOtBu is a stronger oxidant than O₂, the authors suggested that the reaction is initiated by a C–H bond cleavage through the formation of pallada(II)cycle intermediate **3**. The latter is subsequently oxidised by BzOOtBu to give pallada(IV)cycle **4**. The latter intermediate can reductively eliminate a Pd(II) species, forming the C–O bond, and yielding dibenzofuran **2**. As described for the reaction above, also in this case the C–O bond formation occurs at the least sterically hindered position. Notably, substrates bearing either EWGs or EDGs tolerate the oxidative reaction conditions. Building on this work, Schmidt and Riemer²⁰ recently described a microwave-promoted cyclisation protocol, in which oxidative C–H activation using catalytic amounts of Pd(OAc)₂ could be used to prepare DBFs. The reaction works under water-free conditions in non-protic solvents such as benzene and ethereal solvents (THF, MTBE, DME, or DDME).



In 1980, Kappe and co-workers presented a non-CH activated Pd-catalysed diaryl ether formation relying on cyclodehydrogenation of 4-hydroxy-3-phenylquinolinone **5**, quinolinylphenol **7**, and hydroxyphenylphenalenone **9** to prepare the corresponding furan derivatives **6**, **8**, and **10** (Scheme 4).²¹ This method was further used to synthesise biologically active benzofuroquinolinone.²²




Scheme 4 Palladium-catalysed cyclodehydrogenation for the formation of benzofuranes as developed by Kappe and co-workers²¹

2.2 Copper-Mediated C–H Activation


In 2012, Zhu and co-workers published three papers on the dibenzofuran synthesis through C–H bond cleavage/C–O bond formation using Cu-based catalysts.^{23–25} The use of copper is more convenient compared to other transition metals as it is considerably cheaper and compatible with O₂. However, the presence of relatively strong EWGs on the phenolic ring (e.g., nitro, cyano, and carbonyl groups) is required to overcome undesired homocoupling of phenols or other side-reactions usually promoted by copper. In the first protocol, CuBr was used (30%) in the presence of Cs₂-CO₃, and pivalic acid in the air (Scheme 5).²³ Notably, *ortho*-functionalised substrates did not allow formation of the C–O bond, whereas the *meta* congeners gave the cyclised products with a 20:1 regioselective ratio toward the least sterically hindered isomer. For substrates bearing bulky substituents such as *i*-Pr or Ph, Cu(OAc)₂ was used to avoid the formation of undesired brominated by-products. As mentioned above, the phenolic ring can only bear EWGs, whereas the phenyl ring tolerates both EWGs and EDGs.

Together with a KIE value of 4.5, the absence of proton scrambling suggested that the C–H bond cleavage is the rate determining step. As studies with competitive reactions between electron-rich and -poor aryl substrates did not give any conclusive results, the authors suggested that the cyclisation reaction could result from either a concerted metalation-deprotonation (CMD) mechanism or an electrophilic metalation (Scheme 5, paths a,b, respectively). A radical pathway was excluded as the reaction is not affected by the presence of a radical scavenger such as TEMPO. As there are no evidences for the Cu(II)/Cu(0) couple involvement, the oxidative Cu(III)/Cu(I) cycle could not be completely excluded. In a subsequent work,²⁴ the same authors reported on the development of an efficient Cu-catalysed oxidative C–O bond formation of electron-deficient *ortho*-phenols containing supplementary directing *meta*-groups (e.g., NHAc, NHCOPh, 2-pyrrolidone, and NHBoc) on the non-phenolic




Scheme 5 Cu-catalysed DBF formation reported by Zhu and co-workers.²³ Proposed mechanistic pathways: (path a) concerted metalation-deprotonation and (path b) electrophilic metalation.

ring. Together with the hydroxy group, the amidic carbonyl functional group chelates the copper metal centre allowing the reaction to be performed without a base and at low temperatures. Notably, high yields (up to 99 %) and a broad substrate scope with full control on the regioselectivity could be achieved with these substrates (Scheme 6). In a later work, Zhu and co-workers reported the first example of a sequential iodination-cycloetherification reaction of *o*-arylphenols mediated by CuI in the presence of O₂.




Scheme 6 Cu-catalysed DBF formation reported by Zhu and co-workers²⁴

This permitted the synthesis of 2- and 4-iododibenzofurans derivatives using CuI as both iodinating agent and catalyst for the C–H activation/C–O bond formation. Iodination of the phenolic ring at either the *ortho* or *para* positions depending on the position of the EWG was observed (Scheme 7).²⁵ Notably, a thermal control of the iodination reaction to occur prior to the C–O cyclisation could be obtained. Heating the reaction mixture at 60 °C gave first the iodo-inter-



Scheme 7 Cu-catalysed DBF formation reported by Zhu and co-workers²⁵

mediate that, at 140 °C, could be cyclised into the furanyl derivatives in the presence of a stoichiometric amount of CuI (Scheme 8).




Scheme 8 Sequential Cu-catalysed iodination and C–O cyclisation re-ported by Zhu and co-workers.²⁵

This reaction tolerates EWGs (e.g., Cl, F, CF₃, NO₂, CN, and CHO) and EDGs (e.g., Me, OMe, and Ph). Notably, replacing the CuBr with CuI, the corresponding brominated product could be obtained. As in all cases discussed above, the reaction occurs at the least hindered position (regioselectivity 5:1). Capitalising on KIE experiments (4.1) and mechanistic studies using DFT calculations, the authors suggested that the reaction occurs through a pivalate-as-sisted CMD pathway (Scheme 8) initiated by Cu(III) species. Single electron transfer (SET) or electrophilic aromatic substitution (SEAr) mechanisms were excluded.


2.3 Non-CH Activation Oxidant-Mediated Cyclisation

Other convenient synthetic methods to form furan cycles include oxidation methods that do not involve any direct C–H activation, such as (i) transition-metal-free oxidative cyclisations of phenolic ring and trapping of the reactive intermediate by an hydroxyl group²⁶ (Scheme 9), (ii) intramolecular cyclisation of 2,2'-biphenyl-2-quinone²⁷ (Scheme 10), and (iii) oxidation/oxa-Michael cascade reaction (Scheme 11). The latter method relies on the oxidation of a pyrocatechol moiety **31** to the corresponding 1,2-ben-


zoquinone **33**. Subsequent intramolecular 1,4-addition of the neighbouring hydroxyl groups, followed by a tautomerisation/rearomatisation reactions, gave dibenzofuranes **34** and **32**, respectively (Scheme 11). Commonly used oxidants for this transformation are Ag₂O,²⁸ MnO₂,²⁹ and K₃[Fe(CN)₆].³⁰ Capitalising on this reaction, Lu and co-workers prepared derivative **35**, an important intermediate for the total synthesis of (±)-anastatins A and B (Scheme 11).³¹



Scheme 9 Synthesis of asterelin A by Makino et al. involving an oxidative cyclisation²⁶




Scheme 10 Proposed mechanism by Hayashi et al. for the intramolecular cyclisation of tetraphenyl-2,2'-benzoquinone²⁷



Scheme 11 General oxidative formation of DBF through quinone formation (left) as reported by Lu and co-workers³¹

2.4 Light-Mediated Cyclisation


The formation of dibenzofurans was also obtained by photoirradiation of phenols precursors through Excited State Intramolecular Proton Transfer (ESIPT). For example, Wan and co-workers³² observed the formation of diaryl furans in low yield (2–9%) when 1-(2,5-dihydroxyphenyl)naphthalene (**36**) was exposed under irradiation at 300 nm. Notably, different products were obtained depending on the solvent polarity. For instance, in aprotic solvents, a predominant C–C bond migration was observed yielding 2-(2,5-dihydroxyphenyl)naphthalene (**38**) and naphthobenzofuran-8-ol (**37**) as the major (60% yield) and minor (9% yield) products, respectively. In protic solvents, an intramolecular proton transfer followed by electrocyclic ring closure was instead observed, with dihydrobenzoxanthene **40** (53% yield) being the only product (Scheme 12). Molecule **40** can undergo further oxidation to give the pyran derivative.



Scheme 12 Light-induced oxidative formation of DBF motifs³²

2.5 Acid-Catalysed C–O Cleavage/C–O Formation

Biphenol **41** can be transformed into dibenzofuran **2a** through C–O cleavage/C–O formation, likely following a S_EAr mechanism (Scheme 13). This reaction is usually carried out in the presence of either strong Brønsted acids^{33–35} (e.g., PTSA, TfOH, H₂SO₄, and HBr), Lewis acids³⁶ (e.g., SnCl₄, Al₂O₃, and zeolite) or noble metal surfaces [e.g., Au(111) and Ag(111)].³⁷ The latter method has been largely reported in the literature for the preparation of organic materials such as O-doped graphenes³⁷ and helicenes.³⁸



Scheme 13 Formation of DBF from 2,2'-biphenol


3 Six-Membered Rings: DBX, PXX, Xanthone, and Their Derivatives

Concerning the six-membered cyclic ethers, these can be classified as: (i) dibenzoxanthene (DBX), (ii) *peri*-xanthe-noxanthene (PXX), (iii) xanthenes (presenting a carbonyl group joining both aromatic cycles), (iv) miscellaneous derivatives featuring both five- and six-membered rings, and (v) phenoxazines.

3.1 Dibenzoxanthene (DBX)

The first synthetic protocol for the preparation of dibenzoxanthenes (DBX) **43** from a tetra-*tert*-butylated BINOL

derivative was developed in 1963 by Rieche et al.,³⁹ and later optimised by Schneider et al.⁴⁰ Their methods use K₃[Fe(CN)₆] yielding DBXs in 38%. In 2001, Xu et al. discovered the almost quantitative formation of DBX using Cu(II)-amine complexes in hot MeOH in the presence of air.⁴¹ This method tolerates a large variety of amines with the highest reaction rate observed when using ethanolamine. More-over, modification of the alcoholic solvent leads to a variety of derivatives bearing different alkyl ether functionality (Scheme 14). Racemic mixture was obtained when a chiral amine was used or optically pure BINOL was selected as starting material. Based on this observation, they proposed a mechanistic pathway involving the oxidation of BINOL **42** to naphthoxy radical **44**, which is in equilibrium with its carbon-centred radical analogue **45**. The latter radical can be trapped by the alcoholic solvent forming -alkoxy ketone **46**, which could be further oxidised to form DBX **43** (Scheme 14). Later, this synthetic method was used for the



Scheme 14 Oxidative formation of DBX from BINOL⁴¹


synthesis of cancer drugs⁴² or isolated as side product during the deracemisation of BANOL, a phenanthrene derivative of BINOL.⁴³

3.2 Peri-Xanthenoxanthene (PXX)

The first synthesis of PXX dates back to the beginning of the 20th century when Bünzly and Decker⁴⁴ described the oxidation of BINOL in the presence of $K_3[Fe(CN)_6]$. Shortly after, Pummerer used Ag_2O , CuO , and $Cu(OAc)_2$ for the same transformation.⁴⁵ In 2001, Tamotsu and co-workers⁴⁶ revisited Pummerer's protocol, with $Cu(OAc)_2$ in aqueous alkali solution (Table 1).

In 2007, Weinert and co-workers studied the oxidation of 3,3'-disubstituted BINOL in the presence of a sterically encumbered mercury salt, $(Hg[N(SiMe_3)_2])_2$ (**50**).⁴⁷ Depending on the equivalent of oxidant involved in the reaction, they were able to isolate monopyranyl pentacyclic **54**. Based on 1H and ^{199}Hg NMR, they have proposed a mechanistic pathway involving a mercuriation reaction of the hydroxyl groups followed by intramolecular electrophilic aromatic substitution and extrusion of elemental Hg (Scheme 15). Following these reports, Cui and co-workers recently developed an improvement of the synthesis of PXX using $Cu(OAc)_2$ in *ortho*-dichlorobenzene at 190 °C under micro-wave irradiation for only 3 minutes.⁴⁸ In 2016, our group re-reported the synthesis of PXX and PXX-analogues⁴⁹ by adapting Zhu's protocol for dibenzofuran. Specifically, CuI and pivalic acid in DMSO at 140 °C were used. Song and Swager also reported on the electrochemical cyclisation of BINOL derivatives to yield PXX-thiophene-based conducting poly-

mers.⁵⁰ The last to date synthetic protocol for the cyclisation of BINOL to PXX relies on the use of $CuCl$ with *N*-methylimidazole and K_2CO_3 as a base in hot *m*-xylene.⁵¹ This method allows not only the formation of PXX at lower temperature, but to perform cascade dimerisation/cyclisation of simple naphthols. Similarly, Fuchs and co-workers used on-surface protocols to produce DBF- and/or PXX-polymers from 6,6'-dibromo-BINOL derivatives.³⁷



Scheme 15 Proposed formation mechanism for PXX by

Weinert and co-workers⁴⁷

Building on the Cu-based protocols, our group could achieve the synthesis of extended PXX derivatives. In particular, O-doped armchair **56**, **57**⁴⁹ and zig-zag **58**⁵² molecular ribbons, coloured -extended PXX **59** and **60**,³⁵ and monoimide(PXXMI)/diimide(PXXDI) derivatives **61** and **62**

Table 1 Oxidative Formation of PXX from BINOL

Reaction conditions:					
Year	Reagents	R	Solvent	Temp/Time	Yield
1905 ⁴⁴	$[K_3Fe(CN)_6]$	H	—	—	—
1914 ⁴⁵	Ag_2O	H	benzene	80 °C/1 h	—
1914 ⁴⁵	$Cu(OAc)_2/CuO$	H	$PhNO_2$	280 °C/3 h	52–80
2001 ⁴⁶	$Cu(OAc)_2$	H	aq NaOH	r.t./1 h	—
2007 ⁴⁷	$Hg[N(SiMe_3)_2]_2$	H, silyl	benzene	85 °C/24 h	49–97
2013 ⁴⁸	$Cu(OAc)_2$, O_2	<i>n</i> -octyl	ODCB, pyridine	190 °C MWI/3 min	44–56
2016 ⁴⁹	CuI , PivOH, O_2	2,4-di- <i>t</i> BuPh	DMSO	140 °C/2 h	94
2009 ⁵⁰	NOBF ₄	thiophene	CH_2Cl_2	r.t./6 h	42–78
2017 ⁵¹	$CuCl$, NMI, K_2CO_3 , air	H, TMS, alkyl, aryl, CO_2Me , allyl	<i>m</i> -xylene	120 °C/20 h	55–99

were synthesised by our group (Figure 2).⁵³ In-depth photo-physical and electrochemical studies revealed that all pyranopyran derivatives feature strong electron-donating properties due to their high-lying energy HOMO levels.^{52,53} In particular, PXX, PXXMI, and PXXDI revealed to be strong photoreducers, with PXX featuring the same photoreducing potential as that of the commonly used Ir(III) complexes.^{53a}




Figure 2 Chemical structures of extended PXX-based molecular architectures

3.3 Xanthenes

Being xanthenes (i.e., 9*H*-xanthen-9-ones) important building blocks constituting various pharmacological activities, they have been at the centre of a lot of interest in synthetic organic chemistry.⁵⁴ For example, Norathyriol is a chemopreventive agent that is effective against skin cancer.⁵⁵ Daviditin A is used to relax the corpus cavernous smooth muscle, while its Bellidifolin congener is a potent hypoglycemic regulator, improving insulin resistance.⁵⁶ Finally, Atroviridin exhibits anti-inflammatory activity and is traditionally used for the treatment of earache (Figure 3).⁵⁷ In this regard, research groups have focused on the synthetic methodologies that use biogenetic-type approaches to perform the oxidation of the benzophenone core **63**. Polyhydroxylated and methoxylated derivatives were commonly used as substrates and treated with various oxidants such as $K_3Fe(CN)_6$,^{58–61} CrO_3 ,⁵⁹ *p*-chloranil,⁵⁹ $KMnO_4$,^{59,60} $K_2S_2O_8$,⁶⁰ $Mn(OAc)_3$,⁶¹ $Pb(OAc)_4$,⁶¹ and silver salts⁶² (Scheme 16).






Figure 3 Examples of biologically active xanthenes




Scheme 16 Formation of xanthone by oxidation of polyhydroxylated benzophenone

In some cases, enzymatic oxidation,⁶⁰ using Horse Radish Peroxidase Laccase or even microorganism as *R. Buffonii*,⁵⁵ has been used as well. In the case of silver salts,⁶² Fuse et al. suggested a radical mechanism for the formation of 1,7-dihydroxyxanthone **69**. In the proposition, a single electron transfer (SET) to form xanthone ring **67** occurs, generating a radical intermediate stabilised by the 3-hydroxyl group. This is followed by a second SET, resulting in the re-aromatization reaction forming xanthone core **69** (Scheme 17).



Scheme 17 Proposed radical mechanism for the formation of 1,7-dihydroxyxanthone⁶²


Recently, Suzuki et al.⁶³ developed the total synthesis of Atroviridin using MnO_2 as oxidant to form the xanthone core. In their case, the proposed mechanism relies on the formation of *p*-quinone **71** followed by a 1,4-addition of the free hydroxyl group leading to keto-tautomer intermediates **72** and **73**, which are demethylated to yield Atroviridin (Scheme 18).



3.4 Miscellaneous


3.4.1 Conjugated Addition on Quinone

In 2006, Yoshida reported a synthesis of furanyl and pyranyl derivatives based on the conjugated Cu- and Ni-promoted addition of a hydroxyl group on the adjacent *ortho*-quinone moiety (Scheme 19).⁶⁴




During the study of the reaction, they observed that Cu(OAc)₂ was the most effective oxidiser, and that the solvent polarity has a strong effect on the regioselective outcome of the cyclisation. For instance, in the case of DMSO, six-membered ring was preferentially obtained (ratio of 4.4:1), whereas in MeNO₂ only the five-membered ring was formed. Another method for preparing cyclic diaryl ethers is based on Flash Vacuum Pyrolysis (FVP). The idea of this approach is to generate hydroxyl radical species that undergo an intramolecular cyclisation reaction. However, these

reactions are usually low yielding and lack regioselectivity. In their early report, Cardogan and McNab described the cyclisation of 2-(allyloxy) or 2-(benzyloxy)diphenylmethane **81** to produce xanthene **82**, fluorene **83**, and phenol **84** derivative. In the case of 2-(allyloxy)benzophenone **85**, they were able to obtain a mixture of xanthone **86**, fluorenone **87**, methylbenzylphenol **88**, and dibenzofuran **89** (Scheme 20).⁶⁵



3.4.2 Phenoxazine


Phenoxazines (POZs) are natural organic dyes that can be used in applications such as hole-transporting materials, bio-imaging, dye-sensitised solar cells (DSSCs), and lasers.⁶⁶ POZs **92** is usually obtained by oxidative cyclisation of 2-(phenylamino)phenol **91** in the presence MnCl₂,⁶⁷ PbO₂,⁶⁸ or CoCl₂.⁶⁹ Substrates bearing EWGs (e.g., nitro, cyano, and acetyl groups) revealed to be compatible with the oxidative reaction conditions (Scheme 21).





4 Seven-Membered Rings: Cularines

The most common C–H bond cleavage/C–O bond formation developed so far for engineering O-annulated seven-membered rings is that used for the synthesis of cularines. In 1974, Jackson et al. reported an oxidative intramolecular coupling of tetrahydrobenzylisoquinoline derivatives **93a** in the presence of K₃[Fe(CN)₆] to produce cularine **94a** and isocularine **95a** in 2.5%, and 5% yield, respectively.⁷⁰ The

method was improved by using an N-borane complex of **93b**, treating it with VOF₃ to yield **94b** in 56% yield (Scheme 22).⁷¹



Instead, the synthesis of didehydronorcularine **97** was achieved by Rodrigues and Abramovitch using C₆F₅I(OCOCF₃)₂, yielding the desired cyclic diaryl ether in 87%, together with traces of *ortho*-cyclised derivative **98** (Scheme 23).⁷² The same team also reported the synthesis of cularines through the use of nitrenium ions (Scheme 24), generated in situ by acid-catalysed decomposition of the azide precursor **99**. 1,4-Type intramolecular addition of the *peri*-hydroxyl group, followed by tautomerisation of the imine intermediate, yielded amino-didehydronorcularine **102** (81%) as the major product.⁷² Notably, the intramolecular addition reaction occurs at the least sterically hindered site, namely in *para*-position with respect to the amino functionality.



5 Conclusion

In conclusion, in this review we have described the current synthetic approaches to prepare cyclic diaryl ethers through C–H bond cleavage/C–O bond formation. Our attention was focused on five- (furano), six- (pyrano), and seven-membered rings. In the case of the formation of furano-type rings, it is apparent that the C–O bond formation mediated by Pd and Cu salts is generally triggered by the activation of the C–H bond. High yields are usually obtained with metal-catalysed protocols if compared to classical oxidative or light-driven approaches. As far as the pyrano rings are concerned, no clear mechanisms have been postulated so far, and the reaction protocol greatly varies depending on the type of the six-membered diaryl ether, namely, dibenzoxanthene (DBX), *peri*-xanthenoxanthene (PXX), xanthenes, and phenoxazine (POZ). At last, seven-membered diaryl ethers are rare, and only methods to prepare cularines have been discussed.

Funding Information

D. B. thanks the EU through the MC-RISE scheme (INFUSION), and the School of Chemistry @Cardiff University for the financial support.

Acknowledgment

A. R. thanks the FRS-FNRS for his FRIA fellowship. D.B. thanks all present and past co-workers Dr. D. Stassen, Dr. A. Berezin, Dr. A. Sciutto, Dr. D. Biloborodov, Dr. L. Dordevich, Dr. D. Milano, Mr. C. Valentini, and Mr. A. Rossignon for their dedication and achievements in the field of C–O bond formation.

References

- (1) (a) Nicolaou, K. C.; Boddy, C. N.; Bräse, S.; Winssinger, N. *Angew. Chem. Int. Ed.* **1999**, *38*, 2096. (b) Russo, A.; Caggia, S.; Piovano, M.; Garbarino, J.; Cardile, V. *Chem. Biol. Interact.* **2012**, *195*, 1.
- (2) Qu, J.; Xie, C.; Guo, H.; Yu, W.; Lou, H. *Phytochemistry* **2007**, *68*, 1767.
- (3) Teran, T.; Lamon, L.; Marcomini, A. *Atmos. Pollut. Res.* **2012**, *3*, 466.
- (4) (a) Stoessel, P.; Buesing, A.; Heil, H. US Patent 2010/0013381 A1, **2010**. (b) Kobayashi, N.; Sasaki, M.; Ohe, T. US Patent 8 399 288 B2, **2013**. (c) For the *p*-type semiconductor properties of PXX, see: Kobayashi, N.; Sasaki, M.; Nomoto, K. *Chem. Mater.* **2009**, *21*, 552.
- (5) Dou, L.; Chen, C.-C.; Yoshimura, K.; Ohya, K.; Chang, W. H.; Gao, J.; Liu, Y.; Richard, E.; Yang, Y. *Macromolecules* **2013**, *46*, 3384.
- (6) (a) Yang, L.; Li, M.; Song, J.; Zhou, Y.; Bo, Z.; Wang, H. *Adv. Funct. Mater.* **2018**, *28*, 1705927. (b) Li, M.; Guo, Y.; Zhou, Y.; Zhang, J.; Yang, L.; Zhang, L.; Song, J.; Bo, Z.; Wang, H. *ACS Appl. Mater. Interfaces* **2018**, *10*, 13931. (c) Zhou, Y.; Li, M.; Song, J.; Liu, Y.; Zhang, J.; Yang, L.; Zhang, Z.; Bo, Z.; Wang, H. *Nano Energy* **2018**, *45*, 10.
- (7) Lee, C. W.; Seo, J. A.; Gong, M. S.; Lee, J. Y. *Chem. Eur. J.* **2013**, *19*, 1194.

- (7) Wang, Y.; Qiu, S.; Xie, S.; Zhou, L.; Hong, Y.; Chang, J.; Wu, J.; Zeng, Z. *J. Am. Chem. Soc.* **2019**, *141*, 2169.
- (8) Aranyos, A.; Old, D. W.; Kiyomori, A.; Wolfe, J. P.; Sadighi, J. P.; Buchwald, S. L. *J. Am. Chem. Soc.* **1999**, *121*, 4369.
- (9) Ullmann, F.; Sponagel, P. *Ber. Dtsch. Chem. Ges.* **1905**, *38*, 2211.
- (10) (a) Chan, D. M. T.; Monaco, K. L.; Wang, R. P.; Winters, M. P. *Tetrahedron Lett.* **1998**, *39*, 2933. (b) Evans, D. A.; Katz, J. L.; West, T. R. *Tetrahedron Lett.* **1998**, *39*, 2937. (c) Lam, P. Y. S.; Clark, C. G.; Saubern, S.; Adams, J.; Winters, M. P.; Chan, D. M. T.; Combs, A. *Tetrahedron Lett.* **1998**, *39*, 2941.
- (11) Zhu, J. P. *Synlett* **1997**, 133.
- (12) Yi, Z.; Okuda, H.; Koyama, Y.; Seto, R.; Uchida, S.; Sogawa, H.; Kuwata, S.; Takata, T. *Chem. Commun.* **2015**, *51*, 10423.
- (13) Pigge, F.; Coniglio, J. *Curr. Org. Chem.* **2001**, *5*, 757.
- (14) Jalalian, N.; Ishikawa, E. E.; Silva, L. F. Jr.; Olofsson, B. *Org. Lett.* **2011**, *13*, 1552.
- (15) Liu, Z.; Larock, R. C. *Org. Lett.* **2004**, *6*, 99.
- (16) Elbert, S. M.; Reinschmidt, M.; Baumgärtner, K.; Rominger, F.; Mastalerz, M. *Eur. J. Org. Chem.* **2018**, 532.
- (17) (a) Frlan, R.; Kikelj, D. *Synthesis* **2006**, 2271. (b) Pitsinos, E. N.; Vidali, V. P.; Couladouros, E. A. *Eur. J. Org. Chem.* **2011**, 1207. (c) Mandal, S.; Mandal, S.; Ghosh, S. K.; Sar, P.; Ghosh, A.; Saha, R.; Saha, B. *RSC Adv.* **2016**, *6*, 69605. (d) Allen, S. E.; Walvoord, R. R.; Padilla-Salinas, R.; Kozlowski, M. C. *Chem. Rev.* **2013**, *113*, 6234.
- (18) Xiao, B.; Gong, T.-J.; Liu, Z.-J.; Liu, J.-H.; Luo, D.-F.; Xu, J.; Liu, L. *J. Am. Chem. Soc.* **2011**, *133*, 9250.
- (19) Wei, Y.; Yoshikai, N. *Org. Lett.* **2011**, *13*, 5504.
- (20) Schmidt, B.; Riemer, M. J. *Heterocycl. Chem.* **2017**, *54*, 1287.
- (21) Stadlbauer, W.; Schmut, O.; Kappe, T. *Monatsh. Chem.* **1980**, *111*, 1005.
- (22) Chen, Y.-L.; Chung, C.-H.; Chen, I. L.; Chen, P.-H.; Jeng, H.-Y. *Bioorg. Med. Chem.* **2002**, *10*, 2705.
- (23) Zhao, J.; Wang, Y.; He, Y.; Liu, L.; Zhu, Q. *Org. Lett.* **2012**, *14*, 1078.
- (24) Zhao, J.; Wang, Y.; Zhu, Q. *Synthesis* **2012**, *44*, 1551.
- (25) Zhao, J.; Zhang, Q.; Liu, L.; He, Y.; Li, J.; Li, J.; Zhu, Q. *Org. Lett.* **2012**, *14*, 5362.
- (26) Makino, K.; Harada, K.; Kubo, M.; Hioki, H.; Fukuyama, Y. *Nat. Prod. Commun.* **2013**, *8*, 915.
- (27) Hayashi, N.; Kanda, A.; Kamoto, T.; Higuchi, H.; Akita, T. *Heterocycles* **2009**, *79*, 865.
- (28) Aebischer, D.; Brzostowska, E. M.; Mahendran, A.; Greer, A. J. *Org. Chem.* **2007**, *72*, 2951.
- (29) Hirano, M.; Yakabe, S.; Chikamori, H. H.; Clark, J.; Morimoto, T. *J. Chem. Res., Synop.* **1998**, 770.
- (30) Obermeyer, A. C.; Jarman, J. B.; Francis, M. B. *J. Am. Chem. Soc.* **2014**, *136*, 9572.
- (31) (a) Pan, G.; Li, X.; Zhao, L.; Wu, M.; Su, C.; Li, X.; Zhang, Y.; Yu, P.; Teng, Y.; Lu, K. *Eur. J. Med. Chem.* **2017**, *138*, 577. (b) Pan, G.; Ma, Y.; Yang, K.; Zhao, X.; Yang, H.; Yao, Q.; Lu, K.; Zhu, T.; Yu, P. *Tetrahedron Lett.* **2015**, *56*, 4472.
- (32) (a) Nayak, M. K.; Wan, P. *Photochem. Photobiol. Sci.* **2008**, *7*, 1544. (b) Basarić, N.; Došlić, N.; Ivković, J.; Wang, Y.-H.; Veljković, J.; Mlinarić-Majerski, K.; Wan, P. *J. Org. Chem.* **2013**, *78*, 1811.
- (33) (a) Högberg, H.-E.; Komlos, P.; Råmsby, S.; Stjernström, N. E.; Enzell, C. R.; Reid, W. W.; Yanaihara, N.; Yanaihara, C. *Acta Chem. Scand.* **1979**, *33b*, 271. (b) Cui, Y.; Ngo, H. L.; Lin, W. *Inorg. Chem.* **2002**, *41*, 1033. (c) Zhang, C.; Li, T.; Wang, L.; Rao, Y. *Org. Chem. Front.* **2017**, *4*, 386.
- (34) Nakanishi, K.; Fukatsu, D.; Takaishi, K.; Tsuji, T.; Uenaka, K.; Kuramochi, K.; Kawabata, T.; Tsubaki, K. *J. Am. Chem. Soc.* **2014**, *136*, 7101.
- (35) (a) Miletic, T.; Fermi, A.; Orfanos, I.; Avramopoulos, A.; De Leo, F.; Demitri, N.; Bergamini, G.; Ceroni, P.; Papadopoulos, M. G.; Couris, S.; Bonifazi, D. *Chem. Eur. J.* **2017**, *23*, 2363. (b) Papadakis, I.; Bouza, Z.; Stathis, A.; Orfanos, I.; Couris, S.; Miletic, T.; Bonifazi, D. *J. Phys. Chem. A* **2018**, *122*, 5142.
- (36) (a) Petrocelli, F. P.; Klein, M. T. *Ind. Eng. Chem. Prod. Res. Dev.* **1985**, *24*, 635. (b) Arienti, A.; Bigi, F.; Maggi, R.; Moggi, P.; Rastelli, M.; Sartori, G.; Trere, A. *J. Chem. Soc., Perkin Trans. 1* **1997**, 1391. (c) Ogata, T.; Okamoto, I.; Doi, H.; Kotani, E.; Takeya, T. *Tetrahedron Lett.* **2003**, *44*, 2041.
- (37) Kong, H.; Yang, S.; Gao, H.; Timmer, A.; Hill, J. P.; Díaz Arado, O.; Mönig, H.; Huang, X.; Tang, Q.; Ji, Q.; Liu, W.; Fuchs, H. *J. Am. Chem. Soc.* **2017**, *139*, 3669.
- (38) Shyam Sundar, M.; Bedekar, A. V. *Org. Lett.* **2015**, *17*, 5808.
- (39) Rieche, A.; Kirschke, K.; Schulz, M. *Justus Liebigs Ann. Chem.* **1968**, *711*, 103.
- (40) Schneider, H. P.; Streich, E.; Schurr, K.; Pauls, N.; Winter, W.; Rieker, A. *Chem. Ber.* **1984**, *117*, 2660.
- (41) Tan, D.-M.; Li, H.-H.; Wang, B.; Liu, H.-B.; Xu, Z.-L. *Chin. J. Chem.* **2001**, *19*, 91.
- (42) (a) Wang, X.-Z.; Yang, H.-H.; Li, W.; Han, B.-J.; Liu, Y.-J. *New J. Chem.* **2016**, *40*, 5255. (b) Wang, X.-Z.; Yao, J.-H.; Jiang, G.-B.; Wang, J.; Huang, H.-L.; Liu, Y.-J. *Spectrochim. Acta, Part A* **2014**, *133*, 559.
- (43) Hu, G.; Holmes, D.; Gendhar, B. F.; Wulff, W. D. *J. Am. Chem. Soc.* **2009**, *131*, 14355.
- (44) Bünzly, H.; Decker, H. *Ber. Dtsch. Chem. Ges.* **1905**, *38*, 3268.
- (45) Pummerer, R.; Frankfurter, F. *Ber. Dtsch. Chem. Ges.* **1914**, *47*, 1472.
- (46) Takehiro, A.; Norihito, K.; Toshio, N.; Tamotsu, I. *Bull. Chem. Soc. Jpn.* **2001**, *74*, 53.
- (47) Wetherby, A. E. Jr.; Benson, S. D.; Weinert, C. S. *Inorg. Chim. Acta* **2007**, *360*, 1977.
- (48) Lv, N.; Xie, M.; Gu, W.; Ruan, H.; Qiu, S.; Zhou, C.; Cui, Z. *Org. Lett.* **2013**, *15*, 2382.
- (49) Stassen, D.; Demitri, N.; Bonifazi, D. *Angew. Chem. Int. Ed.* **2016**, *55*, 5947.
- (50) Song, C.; Swager, T. M. *Macromolecules* **2009**, *42*, 1472.
- (51) Kamei, T.; Uryu, M.; Shimada, T. *Org. Lett.* **2017**, *19*, 2714.
- (52) Berezin, A.; Biot, N.; Battisti, T.; Bonifazi, D. *Angew. Chem. Int. Ed.* **2018**, *57*, 8942.
- (53) (a) Sciutto, A.; Fermi, A.; Folli, A.; Battisti, T.; Beames, J.; Murphy, D.; Bonifazi, D. *Chem. Eur. J.* **2017**, *24*, 4382. (b) Sciutto, A.; Berezin, A.; Lo Cicero, M.; Miletic, T.; Stopin, A.; Bonifazi, D. *J. Org. Chem.* **2018**, *83*, 13787.
- (54) Pinto, M. M. M.; Sousa, M. E.; Nascimento, M. S. J. *Curr. Med. Chem.* **2005**, *12*, 2517.
- (55) Fromentin, Y.; Grellier, P.; Wansi, J. D.; Lallemand, M.-C.; Buisson, D. *Org. Lett.* **2012**, *14*, 5054.
- (56) Kraus, G.; Mengwasser, J. *Molecules* **2009**, *14*, 2857.
- (57) Tisdale, E. J.; Kochman, D. A.; Theodorakis, E. A. *Tetrahedron Lett.* **2003**, *44*, 3281.
- (58) (a) Atkinson, J. E.; Lewis, J. R. *J. Chem. Soc., Chem. Commun.* **1967**, 803. (b) Graham, R.; Lewis, J. R. *J. Chem. Soc., Perkin Trans. 1* **1978**, 876.
- (59) Ellis, R. C.; Whalley, W. B.; Ball, K. *J. Chem. Soc., Chem. Commun.* **1967**, 803.
- (60) Atkinson, J. E.; Lewis, J. R. *J. Chem. Soc. C* **1969**, 281.
- (61) Shuichi, U.; Kazu, K. *Bull. Chem. Soc. Jpn.* **1977**, *50*, 193.

- (62) Fuse, S.; Matsumura, K.; Johmoto, K.; Uekusa, H.; Tanaka, H.; Hirose, T.; Sunazuka, T.; Ōmura, S.; Takahashi, T. *Chem. Eur. J.* **2016**, *22*, 18450.
- (63) Suzuki, Y.; Fukuta, Y.; Ota, S.; Kamiya, M.; Sato, M. *J. Org. Chem.* **2011**, *76*, 3960.
- (64) Ooyama, Y.; Okamoto, T.; Yamaguchi, T.; Suzuki, T.; Hayashi, A.; Yoshida, K. *Chem. Eur. J.* **2006**, *12*, 7827.
- (65) Cadogan, J. I. G.; Hutchison, H. S.; McNab, H. *J. Chem. Soc., Perkin Trans. 1* **1991**, 385.
- (66) Karlsson, K. M.; Jiang, X.; Eriksson, S. K.; Gabrielsson, E.; Rensmo, H.; Hagfeldt, A.; Sun, L. *Chem. Eur. J.* **2011**, *17*, 6415.
- (67) Bag, S. S.; Ghorai, S.; Jana, S.; Mukherjee, C. *RSC Adv.* **2013**, *3*, 5374.
- (68) Bang, Z. N.; Komissarov, V. N.; Sayapin, Y. A.; Tkachev, V. V.; Shilov, G. V.; Aldoshin, S. M.; Minkin, V. I. *Russ. J. Org. Chem.* **2009**, *45*, 442.
- (69) Lesh, F. D.; Lord, R. L.; Heeg, M. J.; Schlegel, H. B.; Verani, C. N. *Eur. J. Inorg. Chem.* **2012**, 463.
- (70) Jackson, A. H.; Stewart, G. W.; Charnock, G. A.; Martin, J. A. *J. Chem. Soc., Perkin Trans. 1* **1974**, 1911.
- (71) De Lera, A. R.; Saá, J. M.; Suau, R.; Castedo, L. *J. Heterocycl. Chem.* **1987**, *24*, 95.
- (72) Rodrigues, J. A. R.; Abramovitch, R. A.; de Sousa, J. D. F.; Leiva, G. *C. J. Org. Chem.* **2004**, *69*, 2920.